

## **Select BioCycle<sup>®</sup> Plus**



Select BioCycle<sup>®</sup> Plus is a registered trademark of Select Sires Inc. Product listed is manufactured by Agrarian Solutions LLC, Nappanee, IN. Product listed is merely distributed by Select Sires Inc. and manufactured or processed by the company indicated. All claims, representations, and warranties, expressed or implied, are made only by the manufacturer and not by Select Sires Inc.

Updated: January 2025

## A COMPLETE APPROACH TO FEED HYGIENE AND HERD PERFORMANCE CHALLENGES

Defending and protecting cattle against feed hygiene challenges involves two categories of feed additives: binders or probiotic products. Independent, peer-reviewed research indicates that clay and yeast cell wall products are ineffective at "binding" mycotoxins and might actually be more efficacious for binding nutrients, and thereby reducing nutrient bioavailability<sup>1</sup>,<sup>2</sup>.

The Select BioCycle<sup>®</sup> Plus product combines the benefits of Select BioCycle and Select DTX<sup>™</sup> for cattle facing multiple challenges of feed hygiene, digestive pathogens, heat and other factors. The BioCycle Plus direct-fed microbial product features L-form or cell wall-deficient bacteria (CWDB). Independent, peer-reviewed research with the CWDB found improved milk production and reproductive success when lactating dairy cows were challenged by DON and zearalenone<sup>3</sup>,<sup>4</sup>. The BioCycle Plus also features a proprietary blend of natural plant actives sourced from turmeric, naringin and rosemary to defend intestinal integrity and protect liver health for the next level in mycotoxin protection.

USE DIRECTIONS Mix BioCycle Plus into mixed feeds at the rate of: Beef and dairy heifers: 0.25 oz/head/d Dry and lactating cows: Recommended 0.50 oz/head/d

- **PACKAGING** 50 lb. box (#9095)
  - **STABILITY** Stable for 2 years. Store in cool, dry place.

**INGREDIENTS** Montmorillonite clay, calcium carbonate, silicon dioxide, Saccharomyces cerevisiae (active dry yeast), yeast culture, rosemary, sugars, naringin, malic acid, turmeric, potassium iodide, dried egg product, dried Aspergillus oryzae fermentation extract, propionic acid, dried Bacillus subtilis fermentation product, dried Lactobacillus acidophilus fermentation product, tartaric acid, lactic acid, citric acid, aspartic acid.

GUARANTEED	Fungal amylase (A. oryzae)	7,000 SKB/g
ANALYSIS	Fungal protease (A. oryzae)	5,000 HUT/g
	Lipase (Candida cylindracea)	500 FIP/g
	Cellulase (T. reesei)	5,000 CU/g
	Pectinase (A. niger)	100 ENDO-PG/g
	Total microbial activity	
	Bacillus subtilis Lactobacillus acidophilus (min)	60 x 10^6 cfu/g

<sup>1</sup>Kihal et al. (2022). J. Anim. Sci. 100:1-14.
<sup>2</sup>Kihal et al. (2020). J. Dairy Sci. 103:3125-3132.
<sup>3</sup>Roth et al. (2024). J. Dairy Sci. 107(Suppl 1):426 (Abstr.).
<sup>4</sup>Roth et al. (2024). J. Dairy Sci. 107(Suppl 1):426 (Abstr.).







## Select BioCycle<sup>®</sup> Plus Concentrate

Select BioCycle<sup>®</sup> Plus Concentrate is a registered trademark of Select Sires Inc. Product listed is manufactured by Agrarian Solutions LLC, Nappanee, IN. Product listed is merely distributed by Select Sires Inc. and manufactured or processed by the company indicated. All claims, representations, and warranties, expressed or implied, are made only by the manufacturer and not by Select Sires Inc.

Updated: January 2025

## A COMPLETE APPROACH TO FEED HYGIENE AND HERD PERFORMANCE CHALLENGES

Defending and protecting cattle against feed hygiene challenges involves two categories of feed additives: binders or probiotic products. Independent, peer-reviewed research indicates that clay and yeast cell wall products are ineffective at "binding" mycotoxins and might actually be more efficacious for binding nutrients, and thereby reducing nutrient bioavailability<sup>1</sup>,<sup>2</sup>.

The Select BioCycle<sup>®</sup> Plus Concentrate product combines the benefits of Select BioCycle and Select DTX<sup>™</sup> for cattle facing multiple challenges of feed hygiene, digestive pathogens, heat and other factors. The BioCycle Plus Concentrate direct-fed microbial product features L-form or cell wall-deficient bacteria (CWDB). Independent, peer-reviewed research with the CWDB found improved milk production and reproductive success when lactating dairy cows were challenged by DON and zearalenone<sup>3</sup>,<sup>4</sup>. The BioCycle Plus Concentrate also features a proprietary blend of natural plant actives sourced from turmeric, naringin and rosemary to defend intestinal integrity and protect liver health for the next level in mycotoxin protection.

USE DIRECTIONS Mix BioCycle Plus Concentrate into mixed feeds at the rate of: Beef and dairy heifers: 5 g/head/d (0.18 oz) Dry and lactating cows: Recommended 10 g/head/d (0.35 oz)

- PACKAGING 50 lb. bag (#9099)
  - **STABILITY** Stable for 2 years. Store in cool, dry place.

**INGREDIENTS** Montmorillonite clay, calcium carbonate, silicon dioxide, Saccharomyces cerevisiae (active dry yeast), yeast culture, rosemary, sugars, naringin, malic acid, turmeric, potassium iodide, dried egg product, dried Aspergillus oryzae fermentation extract, propionic acid, dried Bacillus subtilis fermentation product, dried Lactobacillus acidophilus fermentation product, tartaric acid, lactic acid, citric acid, aspartic acid.

GUARANTEED	Fungal amylase (A. oryzae)	7,000 SKB/g
ANALYSIS	Fungal protease (A. oryzae)	5,000 HUT/g
	Lipase (Candida cylindracea)	500 FIP/g
	Cellulase (T. reesei)	5,000 CU/g
	Pectinase (A. niger)	100 ENDO-PG/g
	Total microbial activity	
	Bacillus subtilis and Lactobacillus acidophilus	60 million CFU/g

<sup>1</sup>Kihal et al. (2022). J. Anim. Sci. 100:1-14.
<sup>2</sup>Kihal et al. (2020). J. Dairy Sci. 103:3125-3132.
<sup>3</sup>Roth et al. (2024). J. Dairy Sci. 107(Suppl 1):426 (Abstr.).
<sup>4</sup>Roth et al. (2024). J. Dairy Sci. 107(Suppl 1):426 (Abstr.).





www.agrariansolutions.com | Phone (574) 825-1224